Journal of Statistical Planning and Inference 13 (1986) 407–408
North-Holland

AUTHOR INDEX*

Volume 13 (1986)

Baksalary, J.K. and A. Markiewicz, Characterizations of admissible linear estimators in restricted linear models (3) 395–398
Balakrishnan, N. and S. Kocherlakota, On the moments of order statistics from the doubly truncated logistic distribution (1) 117–129
Banerjee, S. and S. Kageyama, A method of constructing regular block designs (3) 399–401
Bartlett, R.F., Estimating the total of a continuous population (1) 51– 66
Beaver, R.J., *see* J.N. Srivastava (2) 133–150
Behnen, K. and G. Neuhaus, Two sample adaptive rank tests under type II censoring (2) 203–213
Bellhouse, D.R. and J.N.K. Rao, On the efficiency of prediction estimators in two-stage sampling (3) 269–281
Bjørnstad, J.F., Asymptotic consistency for subset selection procedures satisfying the P^*-condition (3) 319–335
Borkent, M., *see* N. Cressie (3) 337–344
Buonaccorsi, J.P. and H.K. Iyer, Optimal designs for ratios of linear combinations in the general linear model (3) 345–356
Ceranka, B. and S. Mejza, A note on the inequality $b \geq t$ for block designs (1) 97–101
Chatterjee, K. and R. Mukerjee, Some search designs for symmetric and asymmetric factorials (3) 357–363
Cressie, N. and M. Borkent, The moment generating function has its moments (3) 337–344
Cuesta, J.A. and C. Matrán, L_p-linear regression, consistency and significative regression in median (1) 15– 30
Federer, W.T., *see* D. Raghavarao (2) 151–163
Georgiev, A.A. and W. Greblicki, Nonparametric function recovering from noisy observations (1) 1– 14
Good, I.J., *see* G.I. Holtzman (3) 283–295
Govindarajulu, Z., *see* S.H. Mansouri Ghiassi (2) 239–249
Greblicki, W., *see* A.A. Georgiev (1) 1– 14
Halteman, E., The Chebyshev center: A multidimensional estimate of location (3) 389–394
Hayakawa, T., On testing hypotheses of covariance matrices under an elliptical population (2) 193–202
Holtzman, G.I. and I.J. Good, The Poisson and chi-squared approximations as compared with the true upper-tail probability of Pearson's X^2 for equiprobable multinomials (3) 283–295
Hwang, H.L., On the structure of (v,k,t) trades (2) 179–191
Iyer, H.K., *see* J.P. Buonaccorsi (3) 345–356
Jacroux, M., On the E- and MV-optimality of spring balance designs with string property (1) 89– 96
Kageyama, S., Addendum to 'Connectedness of two-associate PBIB designs' (1) 131–132
Kageyama, S., *see* S. Banerjee (3) 399–401
Kestenband, B.C., Balanced incomplete block designs on $q + 1$ elements (1) 45– 50
Kocherlakota, S., *see* N. Balakrishnan (1) 117–129

* The issue number is given in front of page numbers.

0378-3758/85/$ 3.50 © 1986, Elsevier Science Publishers B.V. (North-Holland)

Kuwada, M., Balanced fractional 2^{m_1} factorial designs of resolution V for interesting effects orthogonal to some effects concerning m_2 factors (3) 365–376
LaRiccia, V.N., Optimal goodness-of-fit tests for normality against skewness and kurtosis alternatives (1) 67– 79
Levy, M.S. and S.K. Perng, A note on the consistency of the modified Fisher prediction function (2) 263–267
Longyear, J.Q., Nested group divisible designs and small nested designs (1) 81– 87
Mansouri Ghiassi, S.H. and Z. Govindarajulu, An asymptotically distribution-free test for ordered alternatives in two-way layouts (2) 239–249
Markiewicz, A., *see* J.K. Baksalary (3) 395–398
Matrán, C., *see* J.A. Cuesta (1) 15– 30
McKean, J.W., *see* G.L. Sievers (2) 215–230
Mejza, S., *see* B. Ceranka (1) 97–101
Mielke, P.W. Jr., Non-metric statistical analyses: Some metric alternatives (3) 377–387
Montakhab, M.S., Embedding of finite pseudo-complements of quadrilaterals (1) 103–110
Mukerjee, R., *see* K. Chatterjee (3) 357–363
Nabeya, S., On a condition for the log-rank test to be a locally optimal rank test (3) 403–405
Nagaraja, H.N., A note on the conditional Markov property of discrete order statistics (1) 37– 43
Nagaraja, H.N., Structure of discrete order statistics (2) 165–177
Neuhaus, G., *see* K. Behnen (2) 203–213
Nishii, R. and T. Shirakura, More precise tables of Srivastava–Chopra balanced optimal 2^m fractional factorial designs of resolution V, $m \leq 6$ (1) 111–116
Perng, S.K., *see* M.S. Levy (2) 263–267
Pilz, J., Minimax linear regression estimation with symmetric parameter restrictions (3) 297–318
Raghavarao, D., W.T. Federer and S.J. Schwager, Characteristics for distinguishing among balanced incomplete block designs with repeated blocks (2) 151–163
Rao, J.N.K., *see* D.R. Bellhouse (3) 269–281
Sathe, Y.S. and R.G. Shenoy, MV-optimal weighing designs (1) 31– 36
Schwager, S.J., *see* D. Raghavarao (2) 151–163
Sen, P.K., *see* J.M. Singer (2) 251–261
Shenoy, R.G., *see* Y.S. Sathe (1) 31– 36
Shirakura, T., *see* R. Nishii (1) 111–116
Sievers, G.L. and J.W. McKean, On the robust rank analysis of linear models with nonsymmetric error distributions (2) 215–230
Singer, J.M. and P.K. Sen, M-methods in growth curve analysis (2) 251–261
Srivastava, J.N. and R.J. Beaver, On the superiority of the nested multidimensional block designs, relative to the classical incomplete block designs (2) 133–150
Wehrhahn, K., Properties of some distributions in randomization theory (2) 231–237